



The Canadian
Continence
Foundation

Continence Facts

PREGNANCY, CHILDBIRTH AND BLADDER CONTROL

Do pregnancy and childbirth affect bladder control?

Yes. But don't panic. If you lose bladder control after childbirth, the problem often goes away by itself. Your muscles may just need time to recover.

When do you need medical help?

If you still have a problem after 6 weeks, talk to your doctor. Without treatment, lost bladder control can become a longterm problem. Accidental leaking can also signal that something else is wrong in your body.

Bladder control problems do not always show up right after childbirth. Some women do not begin to have problems until later, often in their 40's.

You and your health care professional must first find out why you have lost bladder control. Then you can discuss treatment.

After treatment, most women regain or improve their bladder control. Regaining control helps you enjoy a healthier and happier life.

Can you prevent bladder problems?

Yes. Women who exercise certain pelvic muscles have fewer bladder problems later on. These muscles are called pelvic floor muscles. If you plan to have a baby, talk to your doctor. Ask if you should do pelvic floor exercises. Exercises after childbirth also help prevent bladder problems in middle age. Ask your health care professional how to do pelvic exercises.


Constipation, obesity and smoking may also be risk factors.

How does bladder control work?

Your bladder is a muscle shaped like a balloon. While the bladder stores urine, the bladder muscle relaxes. When you go to the bathroom, the bladder muscle tightens to squeeze urine out of the bladder.

More muscles help with bladder control. Two sphincter (SFINK-tur) muscles surround the tube that carries urine from your bladder down to an opening in front of the vagina. The tube is called the urethra (yoo-REE-thrah). Urine leaves your body through this tube. The sphincters keep the urethra closed by squeezing like rubber bands.

Pelvic floor muscles support the bladder and also help keep the urethra closed. When the bladder is full, nerves in your bladder signal the brain. That's when you get the urge to go to the bathroom. Once you reach the toilet, your brain sends a message down to the sphincter and pelvic floor muscles. The brain tells them to relax. The brain signal also tells the bladder muscles to tighten up. That squeezes urine out of the bladder.




The pelvic muscles work to control the release of urine.


(please turn over)



The Canadian Continence Foundation



Healthy bladder
with strong
bladder control
muscles



Leaking bladder
with weak
bladder control
muscles

Strong sphincter (bladder control) muscles prevent urine leakage in pregnancy and after childbirth. You can exercise these muscles to make them strong. Talk to your health care professional about learning how to do pelvic floor exercises.

What do pregnancy and childbirth have to do with bladder control?

The added weight and pressure of pregnancy can weaken pelvic floor muscles. Other aspects of pregnancy and childbirth can also cause problems:


- changed position of bladder and urethra
- vaginal delivery
- episiotomy (the cut in the muscle that makes it easier for the baby to come out)
- damage to bladder control nerves

Which professionals can help you with bladder control?

Professionals who can help you with bladder control include:

- your primary care doctor
- a gynecologist (guy-nuh-CALL-uh-jist): a women's doctor
- a urogynecologist (YOOR-oh-guy-nuh-CALL-uh-jist): an expert in women's bladder problems
- a urologist (yoor-ALL-uh-jist): an expert in bladder problems
- a nurse
- a physiotherapist

Ask if they have an interest in bladder control problems. If not, request that they refer you to a health care professional who specializes in the area. You can also call The Canadian Continence Foundation at 1 800 265-9575 to find the names of those who specialize in bladder control closest to you.



Unborn babies push down on the bladder, urethra and pelvic muscles.

Points to Remember

- Temporary bladder control problems are common during pregnancy.
- Exercising pelvic floor muscles can help prevent or treat bladder control problems.
- Bladder control problems may show up months to years after childbirth. Talk to your health care professional if this happens to you.

For more information about incontinence contact

The Canadian Continence Foundation

P.O. Box 30, Victoria Branch, Westmount, Quebec H3Z 2V4
Tel. (514) 488-8379 ■ Fax (514)488-1379 ■ General information 1-800-265-9575
www.continence-fdn.ca

This reprinting was made possible with an educational grant from Pfizer Canada